

CHANDRA HRC LOCALIZATION OF THE LOW MASS X-RAY BINARIES X1624–490 AND X1702–429: THE INFRARED COUNTERPARTS

STEFANIE WACHTER¹

Spitzer Science Center, California Institute of Technology, MS 220-6, Pasadena, CA 91125

JOSEPH W. WELLHOUSE

Harvey Mudd College, 340 East Foothill Blvd., Claremont, CA 91711

SANDEEP K. PATEL

Universities Space Research Association, NSSTC, SD-50, 320 Sparkman Drive, Huntsville, AL 35805

ALAN P. SMALE²

Laboratory for High Energy Astrophysics/USRA, Code 662, NASA/GSFC, Greenbelt, MD 20771

JOAO F. ALVES

European Southern Observatory, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei München, Germany

PATRICE BOUCHET

Cerro Tololo Inter-American Observatory, National Optical Astronomy Observatory³, Casilla 603, La Serena, Chile
Draft version February 2, 2008

ABSTRACT

We report on the precise localization of the low mass X-ray binaries X1624–490 and X1702–429 with the *Chandra* HRC-I. We determine the best positions to be $16^{\text{h}}28^{\text{m}}02^{\text{s}}825 - 49^{\circ}11'54''.61$ (J2000) and $17^{\text{h}}06^{\text{m}}15^{\text{s}}314 - 43^{\circ}02'08''.69$ (J2000) for X1624–490 and X1702–429, respectively, with the nominal *Chandra* positional uncertainty of $0.6''$. We also obtained deep IR observations of the fields of these sources in an effort to identify the IR counterparts. A single, faint ($Ks = 18.3 \pm 0.1$) source is visible inside the *Chandra* error circle of X1624–490 and we propose this source as its IR counterpart. For X1702–429, a $Ks = 16.5 \pm 0.07$ source is visible at the edge of the *Chandra* error circle. The brightness of both counterpart candidates is comparable to that of other low mass X-ray binary IR counterparts when corrected for extinction and distance.

Subject headings: stars:neutron — X-rays:binaries — X-rays:individual(4U 1624–490, 4U 1702–429)

1. INTRODUCTION

X1624–490 belongs to the group of low mass X-ray binaries (LMXBs) that exhibit periodic intensity dips in their X-ray light curves. It is generally accepted that these so-called “dippers” are high inclination systems and that the dips are due to occultations of the central X-ray source by a thickened region of the accretion disk rim where the gas stream from the companion impacts upon the outer disk. An orbital period of ~ 21 h for X1624–490 was first reported by Watson et al. (1985) and recently refined to 20.89 h by Smale, Church, & Balucińska-Church (2001). X1624–490 is one of the most extreme sys-

tems among the dippers; its persistent X-ray emission is the brightest ($6 \times 10^{37} \text{ erg s}^{-1}$), its dip profiles seemingly the most erratic, and its 20.89-hour orbital period is the longest. The orbital period is ~ 5 –25 times longer than those of the other dipping sources, corresponding to a much greater stellar separation and a larger accretion disk radius. Dipping is deep, $\sim 75\%$ in the 1–10 keV band, and can at times reach a stable lower level (Church & Balucinska-Church 1995) suggesting that one spectral component is entirely removed. The source also exhibits strong flaring in which the X-ray flux can increase by $\sim 30\%$ over timescales of a few thousand seconds. Most recently, Parmar, Oosterbroek, Boirin, & Lumb (2002) discovered narrow absorption features in the *Chandra* X-ray spectrum of X1624–490. Despite the wealth of X-ray observations, the optical/IR counterpart of X1624–490 has not yet been identified. In fact, it is the only one of the known dipping sources without an identified counterpart.

X1702–429 was first discovered as a burst source with OSO 8 (Swank et al. 1976). A persistent X-ray counterpart for this burster was subsequently suggested by Lewin & Clark (1979). This identification

Electronic address: wachter@ipac.caltech.edu
 Electronic address: Sandeep.Patel@msfc.nasa.gov
 Electronic address: alan@osiris.gsfc.nasa.gov
 Electronic address: jalves@eso.org
 Electronic address: pbouchet@noao.edu

¹ also with Eureka Scientific Inc., 2452 Delmer Street, Suite 100, Oakland, CA 94602

² Now at Office of Space Science, NASA Headquarters, Washington DC 20546

³ Operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

was supported by observations from Makishima et al. (1982) who also found variability in the burst activity of the source. Oosterbroek et al. (1991) classified X1702–429 as an Atoll source based on EXOSAT observations. X1702–429 is remarkable because it is one of the few X-ray bursters that shows kHz quasi-periodic oscillations (QPOs) in both its persistent X-ray emission and during type I X-ray bursts, the latter of which might directly indicate the spin period of the neutron star (Strohmayer & Markwardt 1999; Markwardt, Strohmayer, & Swank 1999).

We report here on the precise localization of X1624–490 and X1702–429 with *Chandra* and the subsequent identification of infrared counterpart candidates for both sources.

2. OBSERVATIONS

2.1. X-ray

We observed X1624–490 (Obs ID 2693) on 2002 May 30 and X1702–429 (Obs ID 3791) on 2003 June 19 with the *Chandra* HRC-I for 1 ksec each. CIAO v.3.0 and CALDB v2.26 were used in all data analysis and processing tasks. There were no known aspect corrections for these data sets. We consequently assume an error radius of $0.6''$ (90% confidence) for the derived positions of the X-ray sources from these data according to the information provided at <http://cxc.harvard.edu/cal/ASPECT/celmon/>.

We searched for X-ray sources using both *wavdetect* and our own source finding algorithm. In the X1624–490 data set, a single bright source is detected at the center of the $30' \times 30'$ field. The best position is $16^h28^m02^s.825 -49^\circ11'54''.61$ (J2000) with the $0.6''$ uncertainty mentioned above. The ROSAT All Sky Survey (RASS) sources (1RXS J162622.5-491419 and 1RXS J162824.6-492510) that we hoped to use to improve the global astrometry were not detected. Both are very weak sources in the RASS, and the *Chandra* non-detection implies that they are either not real or transient sources. However, we do find a weak ($0.009 \text{ counts sec}^{-1}$, $S/N \sim 3$) source in our X1624–490 *Chandra* data at $16^h27^m36^s.262 -49^\circ13'03''.31$ (J2000). Its position matches that of the bright 2MASS source 16273627-4913042 at $16^h27^m36^s.271 -49^\circ13'04''.26$ (J2000). The colors and magnitudes of the 2MASS source ($J = 8.936 \pm 0.023$, $H = 8.245 \pm 0.044$, $K = 8.057 \pm 0.024$) are consistent with those of a K4-K5 giant at $\sim 2 \text{ kpc}$ based on the data by Bessell & Brett (1988) transformed to the 2MASS photometric system with the relations by Carpenter (2001). If this association is real, the offset between the *Chandra* and 2MASS astrometric frame is no larger than $0.9''$ and predominantly in the DEC coordinate.

For X1624–490 we determine an HRC-I count rate of $4.25 \pm 0.07 \text{ counts sec}^{-1}$, which falls in the low end of the range predicted using the non-dip, non-flare spectral parameters in the literature (e.g. Church & Balucinska-Church 1995; Bałucińska-Church, Humphrey, Church, & Parmar 2000; Smale, Church, & Bałucińska-Church 2001). According to the source ephemeris (Smale, Church, & Bałucińska-Church 2001), our *Chandra* observation covered phases 0.31–0.34, thus well

away from the intervals where dipping activity would be expected.

In the X1702–429 data set, the X-ray binary is the only source detected, with a count rate of $10.46 \pm 0.10 \text{ counts sec}^{-1}$. This count rate is consistent with the rate we predicted for the *Chandra* HRC-I based on the published flux levels and spectra of this source. Our best localization gives $17^h06^m15^s.314 -43^\circ02'08''.69$ (J2000). Again, the weak RASS source 1RXS J170721.9-431436 we had hoped to use for refining the match between the X-ray and IR astrometric frames is not detected. However, the source would have been at the very edge of the FOV given the roll angle of our *Chandra* observation, so the non-detection is not conclusive.

2.2. Infrared

Observations of the field of X1624–490 were obtained with the CTIO 1.5m and CIRIM on 1997 April 26 - 28 in the *Ks* and *J* filter. A log of all infrared observations of the fields of X1624–490 and X1702–429 is provided in Table 1. The data were dark subtracted and flatfielded in the standard manner using IRAF. A normalized sky was created from the median of the object data, scaled to the median of each image and subtracted. Finally the images were shifted and added into a combined final frame excluding bad pixels via a mask. Photometric and astrometric calibration of this frame was accomplished through a comparison to 2MASS. No IR source is visible in these data at the *Chandra* position of X1624–490 down to a magnitude limit of $Ks \approx 18.0$.

We consequently obtained deeper observations with the ESO NTT and SOFI on 2003 June 19. SOFI was used in the large field mode which provides a $0.29'' \text{ pixel}^{-1}$ scale. The observations were obtained under $\sim 0.8''$ seeing conditions. The data were reduced in a similar manner to the CTIO 1.5m observations described above, with the addition of the recommended crosstalk correction⁴. We derived an astrometric solution for the combined frame through a comparison against 82 2MASS stars, using the IRAF task *ccmap*. The fit has an accuracy of $0.05''$ in each coordinate. Hence, the intrinsic precision of the *Chandra* position is still the dominant source of uncertainty in the localization of the source. We cannot improve the astrometric precision over the nominal $0.6''$ without several other X-ray sources that can be matched to IR sources in the field. We judge the single, weak, additional source detected in our *Chandra* image to be of insufficient quality as a basis for an overall shift in the astrometric solution. Instead, we view it as a large scale check against a significant systematic offset between the two coordinate systems, as mentioned above.

IR observations of the field of X1702–429 were obtained with the CTIO 1.5m and 4m telescopes. Again, the data were reduced in a similar manner to those of X1624–490. As is the case for X1624–490, the CTIO 1.5m data were of insufficient depth to reveal any counterpart candidates for X1702–429. Our most recent CTIO 4m ISPI observations, however, show a counterpart candidate at the edge of the *Chandra* X-ray error circle. ISPI has a pixel scale of $0.3'' \text{ pixel}^{-1}$ and the seeing

⁴ see SOFI Manual, 5.1.4 (August 2003 version), and http://www.eso.org/~gfinger/hawaii_1Kx1K/crosstalk_rock/crosstalk.html

conditions during the observations varied between 1.0–1.2". The astrometric solution for the 4m frame was obtained from comparison to 65 2MASS stars with residuals of 0.04" in each coordinate. We derived photometric calibration transformations for both the X1624–490 ESO and X1702–429 CTIO observations based on a comparison to the 2MASS measurements, neglecting any color terms since we only have data in one filter. The formal rms error in the transformation fit for X1624–490 and X1702–429 was 0.06 and 0.04, respectively.

3. RESULTS

A 1'×1' portion of our combined *Ks* frames as well as a 30"×30" close-up around the *Chandra* position of X1624–490 and X1702–429 are shown in Figure 1 and Figure 2, respectively. The *Chandra* X-ray error circles and several comparison stars are marked. Photometry was performed using *Daophot* (Stetson 1992) and the measurements for all sources indicated in the finding charts for both X1624–490 and X1702–429 are summarized in Table 2. We quote the 2MASS magnitudes for those sources detected with the survey and our own measurements for the fainter sources only resolved with our dedicated observations. Since each potential counterpart is only detected in the combined image of the deepest observations, we are unable to place constraints on the variability of these IR counterparts.

3.1. X1624–490

A single, faint source (X1) is visible inside the *Chandra* error circle, and we propose this source as the IR counterpart to X1624–490. X1 has a magnitude of $Ks = 18.3 \pm 0.1$. X2, the next closest source to the *Chandra* X-ray error circle, is 1.5" away from X1 with a magnitude of $Ks = 18.6 \pm 0.2$. In order to evaluate whether this *Ks* band brightness of X1 is reasonable for a X1624–490 counterpart, we compare it to that of Sco X–1. The two systems have similar orbital periods (Sco X–1: 18.9 h; X1624–490: 20.9 h) and X-ray luminosities (Christian & Swank 1997) and hence might contain a similarly sized accretion disk and mass donor producing the IR emission. One difference between the two systems is the inclination, with Sco X–1 thought to be close to face-on while X1624–490 as a dipper, is close to edge-on. Table 3 lists the extinction corrected absolute *Ks* band magnitudes for both systems, as well as the various measurements that were used in their derivation. We previously determined the brightness of Sco X–1 in *Ks* with dedicated observations and it is also detected with 2MASS. The quoted error in the magnitude for Sco X–1 does not reflect the precision of a single observation but rather the variability of the source when comparing data from multiple epochs. The distance and extinction to Sco X–1 are fairly well known. Bradshaw, Fomalont, & Geldzahler (1999) determined a distance of 2.8 ± 0.3 kpc for Sco X–1 from radio trigonometric parallax. Gallagher, Cash, & Green (1995) claim an upper limit of $E(B - V) = 0.12$ from searches for an X-ray scattering halo, while Willis et al. (1980) favor $E(B - V) = 0.35$ from the observed strength

of the 2200Å band. In any case, due to the reduced effect of optical extinction in the IR ($A_K = 0.114A_V$; Cardelli, Clayton, & Mathis 1989), the extinction at *K* is at most 0.1 mag which does not have a large impact on the derived absolute magnitude. Estimates for the extinction and distance to X1624–490 are much more uncertain. The extinction to X1624–490 is very high, estimated to lie between $A_V = 30 - 50$, based on observed X-ray N_H column densities. A distance estimate of ~ 15 kpc is given by Christian & Swank (1997). The final extinction corrected absolute magnitude range of X1 includes that of Sco X–1 and thus indicates that the observed *K* band brightness is reasonable for that of a counterpart to X1624–490.

3.2. X1702–429

For X1702–429, one IR source (X) is visible at the edge or partially inside the 0.6" *Chandra* error circle. The source appears somewhat elongated and it is not clear whether it would resolve into multiple faint sources at higher resolution. An inspection of the individual IR frames is inconclusive in regards to the morphology of the source. However, *Daophot* has no difficulty detecting and fitting its PSF. Compared to the proposed counterpart for X1624–490, the counterpart candidate for X1702–429 is quite bright at $Ks = 16.5 \pm 0.07$. A comparison to Sco X–1 is not warranted in this case, since X1702–429 is classified as an Atoll source while Sco X–1 is a member of the Z sources. Apart from distinct differences in their X-ray properties, Z and Atoll sources might also indicate particular binary system parameters. Atoll sources generally have shorter orbital periods (3–5 hours) than Z sources and may differ in the type and/or evolutionary state of the mass donor (Bandyopadhyay, Shahbaz, Charles, & Naylor 1999). Unfortunately, the orbital period of X1702–429 is still unknown, so it is difficult to identify an appropriate system for comparison. We have chosen a somewhat typical Atoll source for which we previously obtained IR photometry, the burster X1735–44. X1735–44 has an orbital period of 4.65 h, an estimated extinction of $A_V = 0.7$ and distance of 7–9 kpc (Augusteijn et al. 1998). Table 3 shows that the extinction corrected absolute magnitude of X1702–429 is consistent with that of X1735–44 given the large uncertainties in the input parameters.

The research described in this paper was carried out in part at the Jet Propulsion Laboratory, California Institute of Technology, and was sponsored by NASA *Chandra* grants GO2-3044X and GO3-4036X. This publication makes use of data products from the 2 Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. It also utilized NASA's Astrophysics Data System Abstract Service and the SIMBAD database operated by CDS, Strasbourg, France.

REFERENCES

- Augusteijn, T., van der Hooft, F., de Jong, J. A., van Kerkwijk, M. H., & van Paradijs, J. 1998, *A&A*, 332, 561
 Bałucińska-Church, M., Humphrey, P. J., Church, M. J., & Parmar, A. N. 2000, *A&A*, 360, 583

TABLE 1
LOG OF IR OBSERVATIONS

Date	Telescope + Instr.	Filter	# Coadds $\times$ Exptime (s)	# of Exp.
X1624–490				
1997 April 28	CTIO 1.5m + CIRIM	<i>J</i>	4×25	27
1997 April 26–28	CTIO 1.5m + CIRIM	<i>Ks</i>	4×25	151
2003 June 19	ESO NTT + SOFI	<i>Ks</i>	10×5	17
X1702–429				
1995 June 03	CTIO 1.5m + CIRIM	<i>Ks</i>	4×15	9
2004 July 24	CTIO 4m + ISPI	<i>Ks</i>	3×20	9

TABLE 2
IR PHOTOMETRY

Source	<i>Ks</i>	2MASS Designation
X1624–490		
A	10.623 ± 0.025	16280335-4911535
B	12.572 ± 0.025	16280420-4911552
C	14.81 ± 0.07	
X1	18.31 ± 0.1	
X2	18.63 ± 0.2	
X1702–429		
A	12.081 ± 0.026	17061561-4302042
B	16.17 ± 0.07	
C	12.181 ± 0.026	17061714-4301492
X	16.47 ± 0.07	

TABLE 3
RESULTS

Source	<i>Chandra</i> Coordinates (J2000)	m_{Ks}	A_V	Distance (kpc)	M_{Ks0}	Ref.
X1624–490	16:28:02.825 –49:11:54.61	18.31 ± 0.1	30–50	15	–0.99 to –3.27	1
Sco X–1	...	11.2 ± 0.3	0.3–0.9	2.8 ± 0.3	–1.14	2,3,4
X1702–429	17:06:15.314 –43:02:08.69	16.47 ± 0.07	5	7	1.67	5
X1735–44	...	16.77 ± 0.2	0.7	7–9	2.46–1.92	6,1

REFERENCES. — (1)Christian & Swank (1997) (2)Gallagher, Cash, & Green (1995) (3)Willis et al. (1980)
(4)Bradshaw, Fomalont, & Geldzahler (1999) (5)Oosterbroek et al. (1991) (6)Augusteijn et al. (1998)

Bandyopadhyay, R. M., Shahbaz, T., Charles, P. A., & Naylor, T. 1999, MNRAS, 306, 417
 Bessell, M. S., Brett, J. M. 1988, PASP, 100, 1134
 Bradshaw, C. F., Fomalont, E. B., & Geldzahler, B. J. 1999, ApJ, 512, L121
 Cardelli, J. A., Clayton, G. C., Mathis, J. S. 1989, ApJ, 345, 245
 Carpenter, J. M. 2001, AJ, 121, 2851
 Christian, D. J. & Swank, J. H. 1997, ApJS, 109, 177
 Church, M. J. & Balucinska-Church, M. 1995, A&A, 300, 441
 Gallagher, D., Cash, W., & Green, J. 1995, ApJ, 439, 976
 Landolt, A. U. 1992, AJ, 104, 340
 Lewin, W. H. G. & Clark, G. W. 1979, Proc. Symposium on the Results and Future Prospects of X-ray Astronomy (Tokyo), p.3
 Makishima, K., et al. 1982, ApJ, 255, L49
 Markwardt, C. B., Strohmayer, T. E., & Swank, J. H. 1999, ApJ, 512, L125
 Oosterbroek, T., Penninx, W., van der Klis, M., van Paradijs, J., & Lewin, W. H. G. 1991, A&A, 250, 389

Parmar, A. N., Oosterbroek, T., Boirin, L., & Lumb, D. 2002, A&A, 386, 910
 Predehl, P., & Schmitt, J. H. M. M., 1995, A&A, 293, 889
 Smale, A. P., Church, M. J., & Bałucińska-Church, M. 2001, ApJ, 550, 962
 Stetson, P. B. 1992, ASP Conf. Ser. 25: Astronomical Data Analysis Software and Systems I, 1, 297
 Strohmayer, T. E. & Markwardt, C. B. 1999, ApJ, 516, L81
 Swank, J. H., Becker, R. H., Pravdo, S. H., Saba, J. R., & Serlemitsos, P. J. 1976, IAU Circ., 3010, 1
 Watson, M. G., Willingale, R., King, A. R., Grindlay, J. E., & Halpern, J. 1985, IAU Circ., 4051, 2
 Willis, A. J., et al. 1980, ApJ, 237, 596




FIG. 1.— *Ks* band image of the field of X1624–490, a $1' \times 1'$ field of view on the left and a $30'' \times 30''$ close-up on the right. The best *Chandra* position of $16^h 28^m 02^s.825 - 49^\circ 11' 54''.61$ (J2000) with an error circle of $0.6''$ radius is shown. A faint source (X1) is visible inside the error circle.




FIG. 2.— *Ks* band image of the field of X1702–429, a $1' \times 1'$ field of view on the left and a $30'' \times 30''$ close-up on the right. The best *Chandra* position of $17^h 06^m 15^s.314 - 43^\circ 02' 08''.69$ (J2000) with an error circle of $0.6''$ radius is shown. A faint source (X) is visible at the edge of the error circle.